



Center for Eukaryotic Structural Genomics

Protein Structure Data Summary

Target ID	GO.15838	
Source Organism	<i>Arabidopsis thaliana</i>	
Target Name	At3g25760.1	
PDB Entry	1ZVC	Deposition: 01-Jun-2005
Function	Allene oxide cyclase	
Produced From	<i>E. coli</i> B834 p(RARE2) pVP-16	
Structure by X-ray	Resolution: 1.79 Å	R-value (R-free): 23.9% (28.3%)
	No. of Residues/ASU: 171	Monomers/ASU: 1
Data Collected At	Advanced Photon Source 22-BM 12-Mar-2005	
Authors	G.E. Wesenberg, G.N. Phillips, Jr., B.W. Han, E. Bitto, C.A. Bingman, S.T.M. Allard	



Structural Features

Arabidopsis thaliana gene At3g25770.1 encodes an isoform of allene oxide cyclase (AOC1), a key enzyme involved in biosynthesis of jasmonates. Jasmonates are plant hormones with growth-inhibiting and senescence-promoting activities. They also act as signaling molecules in plant defense by inducing the expression of protease inhibitors and antimicrobial agents. The biosynthesis of jasmonates starts with linolenic acid. Allene oxide cyclase catalyzes the cyclization of an unstable epoxide precursor of jasmonate, generating the first biologically active compound of the pathway, 12-oxo-phytodienoic acid (OPDA). The crystal structure of AOC1 reveals the same fold as AOC2, another AOC isoform solved by CESH (PDB ID 1Z8K). The two isoforms superimpose very well with RMSD of 0.30 Å for 162 atoms. Sequence comparison of AOC1 and AOC2 shows total of ten mutations. Most of them are conservative changes on the protein surface with no differences in ligand binding site. This is in agreement with the results of an *in vitro* assay that showed similar activity and substrate specificity for all four AOC isoforms found in *A. thaliana* (1). However, differences in expression pattern point toward a specific function of each AOC isoform. AOC2 is upregulated upon wounding whereas AOC1 is strongly expressed during senescence. Small structural differences between AOC1 and AOC2 might facilitate the interaction with different regulatory proteins resulting in functional specificity of those two proteins.

References: (1) Hofmann, E., Zerbe, P., Schaller, F. (2006) The crystal structure of *Arabidopsis thaliana* allene oxide cyclase: insights into the oxylipin cyclization reaction. *Plant Cell* 18:3201-3217.

Percent Identity with Nearest PDB Structure at Time Solved	93% (1Z8K)
Pfam Cluster	Allene_oxide_cyclase
Sequence Cluster Size	36

Center for Eukaryotic Structural Genomics (CESG), University of Wisconsin-Madison Biochemistry Department, 433 Babcock Drive, Madison, WI 53706-1549; phone: 608.263.2183; fax: 608.890.1942; email: cesginfo@biochem.wisc.edu; website: <http://www.uwstructuralgenomics.org>. This research funded by NIH / NIGMS Protein Structure Initiative grants U54 GM074901 and P50 GM064598.