

Center for Eukaryotic Structural Genomics


Technology Dissemination Report

CESG Tech Report No.	030
Title	Detergent Solubilization Screening for Membrane Proteins Expressed in <i>E.coli</i>
Research Unit	Protein Production (Large Scale Protein Production)
Authors	Gromek, K.A., Beebe, E.T., Frederick, R.O., Bergeman, L., Hwang S., Li, H., Nichols, K.W., Aceti, D.J., Vojtik, F., Primm, J.G., Ruoho, A.E., Chu, U. B., Markley, J.L., Phillips, G.N., Jr., and Fox, B.G.

FC-12 →	0%	0.031%	0.05%	0.07%	0.10%	0.20%	0.50%	0.70%	1%
Triton X-100 ↓	1	2	3	4	5	6	7	8	9
A 0.01%	0% FC-12 0.01% Triton X-100	0.031% FC-12 0.01% Triton X-100	0.05% FC-12 0.01% Triton X-100	0.05% FC-12 0.01% Triton X-100	0.1% FC-12 0.01% Triton X-100	0.2% FC-12 0.01% Triton X-100	0.5% FC-12 0.01% Triton X-100	0.7% FC-12 0.01% Triton X-100	1% FC-12 0.01% Triton X-100
B 0.05%	0% FC-12 0.05% Triton X-100	0.031% FC-12 0.05% Triton X-100	0.05% FC-12 0.05% Triton X-100	0.05% FC-12 0.05% Triton X-100	0.1% FC-12 0.05% Triton X-100	0.2% FC-12 0.05% Triton X-100	0.5% FC-12 0.05% Triton X-100	0.7% FC-12 0.05% Triton X-100	1% FC-12 0.05% Triton X-100
C 0.1%	0% FC-12 0.1% Triton X-100	0.031% FC-12 0.1% Triton X-100	0.05% FC-12 0.1% Triton X-100	0.05% FC-12 0.1% Triton X-100	0.1% FC-12 0.1% Triton X-100	0.2% FC-12 0.1% Triton X-100	0.5% FC-12 0.1% Triton X-100	0.7% FC-12 0.1% Triton X-100	1% FC-12 0.1% Triton X-100
D 0.5%	0% FC-12 0.5% Triton X-100	0.031% FC-12 0.5% Triton X-100	0.05% FC-12 0.5% Triton X-100	0.05% FC-12 0.5% Triton X-100	0.1% FC-12 0.5% Triton X-100	0.2% FC-12 0.5% Triton X-100	0.5% FC-12 0.5% Triton X-100	0.7% FC-12 0.5% Triton X-100	1% FC-12 0.5% Triton X-100
E									
F									
G 0%	0% FC-12 0% Triton X-100	0.031% FC-12 0% Triton X-100	0.05% FC-12 0% Triton X-100	0.05% FC-12 0% Triton X-100	0.1% FC-12 0% Triton X-100	0.2% FC-12 0% Triton X-100	0.5% FC-12 0% Triton X-100	0.7% FC-12 0% Triton X-100	1% FC-12 0% Triton X-100

Solubilization array for MBP-human sigma1 receptor. The concentrations of detergents that solubilize over 90% of protein are marked in yellow.

Solubilization screening of MBP-sigma1 receptor (LA.25752) in presence of (A) Triton X-100 and fos-choline-12 (FC-12) and (B) fos-choline-12 (FC-12). S- soluble fraction, P- insoluble fraction (pellet).



Summary

CESG is developing new purification methods for membrane proteins expressed in *Escherichia coli* (*E. coli*), that is based on a high-throughput detergent solubilization screen and IMAC protein purification. In case there is no previous research done on a particular membrane protein, this screening procedure starts with wide variety of detergents across a range of concentrations.

Here we present an approach to solubilize a human sigma1 receptor expressed as N-terminal MBP fusion (with signal sequence that directs it to the periplasm – plasmid pVP87K). This protein has been extensively studied but the structure has yet to be solved [1]. To perform this screen in a 96 well plate, it is sufficient to start with 0.1 g of wet cell paste. After breaking the cells by sonication, the lysate is centrifuged at 16,000g to separate the insoluble and soluble fractions. The pellet (insoluble fraction) is then re-suspended and used in the high-throughput detergent solubilization screens. The testing is performed at 4°C overnight, and then the solubilized fraction is separated by centrifugation. We employ the Bio-Rad stain free imager to estimate the efficiency of solubilization.

This method can be semi- or fully automated, and is serving as a precursor for automated purification using the Maxwell 16 system.

[1] Ramachandran, S., Lu, H., Prabhu, U. and. Ruoho, A.E. (2007) Purification and characterization of the guinea pig sigma-1 receptor functionally expressed in *Escherichia coli*. *Protein Expr Purif.* 51(2): 283-92'

Acquiring the Technology

Contact John Primm jgprimm@wisc.edu

Other Acknowledgements

Center for Eukaryotic Structural Genomics (CESG), University of Wisconsin-Madison Biochemistry Department, 433 Babcock Drive, Madison, WI 53706-1549; phone: 608.263.2183; fax: 608.890.1942; email: cesginfo@biochem.wisc.edu; website: <http://www.uwstructuralgenomics.org>. This research funded by NIH / NIGMS Protein Structure Initiative grant U54 GM074901